EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Teldor (PCS No. 02003)

CROPS:

[Field & Protected Crops of: lettuce, herbs, baby leaf brassica, parsley & Protected crops of courgettes, cucumber, pepper, aubergine, gherkin and tomato]

For this EAMU the following crop definitions apply for herbs, lettuce and baby leaf:

Basil, borage, caraway, chervil, coriander, cress, curry plant, dill, fenugreek, feverfew, frise, lambs lettuce, lettuce, marigold, nasturtium, parsley, rocket (salad) savory, scarole, violet, *Agastache spp*, angelica, balm, bay, burnet (salad) catnip, chamomile, chive, clary, sage, dragonhead, fennel as a leafy herb, horehound, hyssop, land cress, lavandin, lavender, lemon verbena, lovage, marjoram, mint, nettle, oregano. *Origanum heracleoticum*, plantain, rosemary, rue, sage, sorrel, spike lavender, St John's Wort, tarragon, thyme, woodruff, yarrow

This approval provides for the use of the product <u>Teldor (PCS No. 02003)</u> in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor use. The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 31.01.2018. [Includes latest changes. This version should be used until updated]

Product name: Teldor

Active Substance: Fenhexamid 51% w/w

Formulation: Water Dispersible Granule

Authorisation Holder: Bayer CropScience Limited

Marketing Company: Bayer CropScience Limited

PCS No: 02003

User type/Function: Professional Use / Fungicide.

Application:

Crop	Max. individual dose	Max. number of applicatio ns	Max. total dose	Minimum interval between applications	time of application	Target
(Field and protected) Lettuce, herbs, baby leaf brassica and parsley	1.5 kg/ha	2 per season	3 kg/ha	-	3 days pre harvest	Botrytis cinerea (grey mould)
(Protected) Courgette, cucumber, pepper, aubergine, gherkin and tomato	1.5 kg/ha*	3 per season	4.5 kg/ha	-	1 day pre harvest	Botrytis cinerea (grey mould)

^{*} Applications to be made by hand-held sprayer or gantry sprayer at a maximum concentration of 100grams of product per 100litres of water & a maximum individual dose of 1.5kg of product in 1500litres of water per ha. Spray to run off.

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed sprayer, gantry sprayer and knapsack/handheld sprayers only.

- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Teldor as an agricultural fungicide for use on the crops and situations listed above.

Applications should be applied protectantly or at the first signs of disease using a minimum of 600 litres of water per ha (1500 litres of water per ha for crops of aubergine, courgette, cucumber, gherkin, pepper & tomato. Growers are advised to test a small area of the crop first to ensure the application will not cause damage to the crop especially if the crop is under any stress.

Resistance Management:

This product should be applied in accordance with an anti-resistance strategy. Disease control may be reduced if strains of the pathogen less sensitive to Fenhexamid develop.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical types or alternative control measures should be included in the planned programme. Alternating fungicides with different modes of action is a recognised anti-resistance strategy and Teldor should always be used in conjunction with other fungicides with a different mode of action / different cross resistance group with satisfactory efficacy against the target.

Fenhexamid is classed (FRAC 17) & is an SBI:Class III hydroxyanilide fungicide.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.