

**EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)****INSTRUCTIONS FOR USE****PPP: SL567A (PCS No. 03106)**

CROPS:

[Field: Root and tuber vegetables and Brassica vegetables.]

[Protected: Tomato]

**For this EAMU the following crop definitions apply and include:**

**Field crops of -**

**Root and tuber vegetables include:**

Parsnip, radish, horseradish

**Brassica vegetables include:**

Brussels Sprout, head cabbage & cauliflower.

This approval provides for the use of the product **SL567A (PCS No. 03106)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

**Reference Date: 09.10.20.** [Includes latest changes. This version should be used until updated]

*Product name:* SL567A

*Active Substance:* Metalaxyl M

***There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.***

*Formulation:* 465.2g/L SL  
*Authorisation Holder:* Syngenta UK Ltd  
*Marketing Company:* Syngenta Ireland Ltd  
*PCS No:* 03106  
*User type/Function:* Professional Use Only / Fungicide.  
*Application:*

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Field: parsnip, radish & Horseradish]	1.3 L/ha	1 per crop	1.3 L/ha	n/a	No later than 6 weeks after drilling and 30 days before harvest.	Cavity spot caused by <i>Pythium violae</i> .
[Protected: Tomato]	0.3 L/ha	3 per crop	0.9 L/ha	3 days	3 days before harvest	Botrytis & <i>phytophthora infestans</i> .
Field: Brussels sprout, Head cabbage & cauliflower.	0.16 L/ha	2 per crop	0.32 L/ha/crop	14 days	14 days before harvest	White blister ( <i>Albugo candida</i> )

*Other specific restrictions:*

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application may be made by tractor mounted/trailed/air assisted sprayer **only** for the field crops listed & via overhead gantry and handheld sprayer **only** for the protected crops.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.

***There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.***

- (5) For all the crops listed this product must be applied in a minimum of 200 litres of water per ha.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above. Specifically, for the indoor application: operators MUST be fully protected [covered] including suitable respiration equipment when making the application.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

*Advisory information:*

This approval relates to use of SL567A as a horticultural fungicide for the control; of Cavity spot on the root crops listed, botrytis plus other fungal diseases [Phytophthora infestans] on tomato & white blister on the brassica crops listed.

The root crops may be applied between 200 and 1000 litres of water per ha.

Tomato crops should be sprayed to run off.

Brassica crops to run off but in a minimum of 200 litres.

SL567A is a fungicide with systemic and protectant activity against pythium violae but has no control on pythium sulcatum (one of the lesser causative agents) of cavity spot.

*Resistance Management:*

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Metalaxyl-M, a component of SL 567A and belongs to the chemical class phenylamides. A number of diseases have developed resistance to phenylamides. Where strains of pathogens resistant to metalaxyl-M develop SL 567A may not give satisfactory control of the disease. Where there is a known resistance problem with metalaxyl-M growers should seek other methods of disease control.

To prevent the build-up of resistance which could render SL-567A ineffective, when applied to foliage it should be used in mixture with fungicides having a different mode of action. Cultural methods of control should be used whenever possible. SL-

***There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.***

567A should only be used where essential. Curative or eradicant use should be avoided. To minimise the risk of resistance developing this product must always be applied in tank-mix with another fungicide with an alternative mode of action and at a rate that is effective against the target disease. There is a high risk of resistance developing if SL 567A is applied alone. SL567A contains metalaxyl M which is classed by FRAC as group: 4 - Phenylamides.

*IMPORTANT PLEASE NOTE:*

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

---

***There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.***