

**EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)****INSTRUCTIONS FOR USE****PPP: Sparviero (PCS No. 04469)**

CROPS: Outdoor crops of: Baby leaf\*, beetroot, celeriac, celery, herbs\*\*, horseradish, Jerusalem artichoke, parsley root, radish, salsify, spinach, swede, turnip.

For this EAMU the following crop definitions apply:

\*Baby leaf production includes leaves of lettuces, spinach and leafy brassicas harvested up to 8 true leaves.

\*\* Perennial herbs {*Agastache spp*: angelica, balm, bay, burnet (salad), catnip, chamomile, chives, clary sage, dragonhead, fennel, horehound, hyssop, land cress, lavandin, lavender, lemon verbena, lovage, marjoram, mint, nettle, oregano}, {*Origanum heracleoticum spp*: plantain, rosemary, rue, sage, sorrel, spike lavender, St john's Wort, tarragon, thyme, woodruff, yarrow}

Annual herbs: {basil, Borage, caraway, chervil, coriander, curry plant, dill, fenugreek, feverfew, marigold, nasturtium, parsley, rocket (Rucolla), savory, violet}

This approval provides for the use of the product **Sparviero (PCS No. 04469)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

**Reference Date: 25.02.19.** [Includes latest changes. This version should be used until updated]

***There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.***

*Product name:* Sparviero

*Active Substance:* Lambda-cyhalothrin

*Formulation:* 100g/l CS

*Authorisation Holder:* Sipcam Oxon S.p.A.

*Marketing Company:* Sipcam UK Ltd.

*PCS No:* 04469

*User type/Function:* Professional Use Only / Insecticide.

***Application:***

| Crop   | Max. individual dose          | Max. number of applications | Max. total dose | Min. interval between applications | Latest time of application | Target                                 |
|--|-------------------------------|-----------------------------|-----------------|------------------------------------|----------------------------|--|
| Field baby leaf production (up to 8 true leaves) & field herbs                                     | 75 ml/ha (min 400L water/ha)  | 4                           | 150 ml/ha       | 10 days                            | 7 days pre-harvest         | Cutworms                               |
| Field spinach  | 75 ml/ha (min 400L water/ha)  | 4                           | 150 ml/ha       | 10 days                            | 7 days pre-harvest         | Cutworms                               |
| Field celery   | 50 ml/ha (min 300L water/ha)  | 4                           | 150 ml/ha       |                                    | 7 days pre-harvest         | Silver y moth caterpillars, Carrot fly |
| Beetroot, horseradish, jerusalem artichoke, celeriac, parsley root, radish, salsify, swede, turnip | 150 ml/ha (min 300L water/ha) | 4                           | 450 ml/ha       |                                    | 14 days pre-harvest        | Carrot fly, cabbage fly                |

***There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.***

*Other specific restrictions:*

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application may only be made using a tractor mounted sprayer, knapsack, hydraulic nozzle applicator.
- (4) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.
- (5) This approval extends to professional use only.
- (6) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5 metres to surface water bodies from horizontal boom sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (7) For operator protection refer to the product label.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

*Advisory information:*

This approval relates to the use of Sparviero as an agricultural/horticultural insecticide for the crops & situations listed above.

**Resistance Management:**

Use Sparviero as part of an Integrated Crop Management (ICM) strategy.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other insecticides from different cross-resistant groups [different mode of action, IRAC group] with satisfactory efficacy against the target.

Sparviero contains Lambda-cyhalothrin IRAC class: 3A.

**IMPORTANT PLEASE NOTE:**

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

---

***There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.***