

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)**INSTRUCTIONS FOR USE****PPP: Volley (PCS No. 04516)****CROPS:**

[Outdoor grown: Carrot, celeriac, parsley (root) and beetroot]

This approval provides for the use of the product **Volley (PCS No. 04516)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 10.05.22. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Volley
<i>Active Substance:</i>	Fluazinam
<i>Formulation:</i>	500 g/L Suspension Concentrate (SC)
<i>Authorisation Holder:</i>	Adama Agricultural Solutions UK Ltd
<i>Marketing Company:</i>	Adama Agricultural Solutions UK Ltd
<i>PCS No:</i>	04516
<i>User type/Function:</i>	Professional Use Only / Fungicide

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Carrot & Celeriac	0.2 l/ha	2	0.4 l/ha	7-10 days	7 days before harvest	Alternaria
Parsley (root)	0.2 l/ha	2	0.4 l/ha	7-10 days	7 days before harvest	Alternaria
Beetroot	0.4 l/ha	1	0.4 l/ha	-	7 days before harvest	Alternaria alternate

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed horizontal sprayer only.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label and specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed horizontal sprayers. Direct spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Volley as a horticultural fungicide for use on the crops and situations listed above.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Applications should be applied protectantly or at the first signs of disease using a minimum of 200 litres of water per ha for red beet and 700 litres of water for carrots, parsley (root) and celeriac.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in rotation & in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Volley contains Fluazinam

Fluazinam is classed FRAC group: uncouplers of oxidative phos-phorylation.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.