

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Gamit 36 CS (PCS No. 05357)

CROPS:

Outdoor grown: Cabbage, cauliflower, broccoli, brussels sprout, calabrese, Chinese cabbage, collards, kale, choi sum, pak choi, courgette, carrot (post emergence use) and swede.

This approval provides for the use of the product **Gamit 36 CS (PCS No. 05357)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 13.07.18. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Gamit 36 CS
<i>Active Substance:</i>	Clomazone
<i>Formulation:</i>	360 g/l capsule suspension
<i>Authorisation Holder:</i>	FMC Chemical s.p.r.l
<i>Marketing Company:</i>	Headland Agrochemicals Ltd
<i>PCS No:</i>	05357
<i>User type/Function:</i>	Professional Use Only / herbicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Min. interval between applications	Latest time of application	Target
Outdoor grown: cabbage, cauliflower, broccoli, brussels sprout, calabrese, Chinese cabbage, collards, kale, choi sum, pak choi	0.25 l/ha	-	0.25 l/ha	-	Before 6 true leaf stage (early post transplantation)	See below
Outdoor courgette	0.25 l/ha	-	0.25 l/ha	-	Before 2 nd true leaf stage	See below
Outdoor swedes	0.25 l/ha	-	0.25 l/ha	-	Pre-emergence	See below
Outdoor carrot (Post emergence use)	83 ml/ha	3 per crop	0.25 l/ha	14 days	BBCH 14 (4 true leaves) & 55 days before harvest.	See below

Target Weeds:

Groundsel, cleavers, fools parsley, chickweed, red dead nettle, redshank, ivy leaved speedwell and wild radish

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient. This specific restriction also includes any application applied pre-emergence. That is, when using on carrots the total dose of Gamit 36 CS that may be applied pre and post emergence is 0.250 l/ha/crop.
- (3) Application by tractor mounted/trailed sprayer only.
- (4) Applications of any products containing clomazone must not be applied to two or more consecutive crops of courgette in one calendar year.
- (5) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (6) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 1m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (7) This approval extends to professional use only.
- (8) This product may only be used where there is a minimum re-planting/sowing interval of 6 weeks for any new crop
- (9) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (10) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Gamit 36 CS as a horticultural herbicide for the crops and situations listed above.

Apply in a minimum of 200 litres of water per Ha.

Application can be made to drilled crops from pre-emergence up to the stated latest time of application. Transplanted crops should be allowed to harden off before treatment.

Safety Note --- Drilled cabbage --- only apply Gamit 36 CS post-emergence of drilled cabbage when at minimum 2 true leaf growth stage.

Growers are cautioned that whilst there is some evidence that suggests a possible safe use in certain soils and conditions, there has been no complete testing of this product post emergence in all situations. Growers are strongly advised to test a small area before widespread use. Some of the factors that may effect the phytotoxicity of this product include, temperature, soil moisture content, soil sand component, leaf wax etc. Growers are reminded that this EAMU is commercially at the growers own risk. Please consult with your crop advisers for more specialist advice.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other herbicides from different cross-resistant groups [different mode of action, HRAC group] with satisfactory efficacy against the target.

Gamit 36 CS contains clomazone

Clomazone is classed HRAC group: F3.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.