

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)**INSTRUCTIONS FOR USE****PPP: Switch (PCS No. 06414)**

CROPS: Protected crops of: blueberry, cucumber, gherkin, courgette, tomato, pepper & aubergine.

Outdoor crops of: asparagus, bulb onion, garlic, shallot, lupins, salad onion, beetroot, horseradish, parsnip, parsley root, salsify, cherry, plum & celery.

Outdoor and protected crops of: cress, endives, Lambs lettuce, lettuce, rocket, herbs, spinach, spinach beet & baby leaf brassica

This approval provides for the use of the product **Switch (PCS No. 06414)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 05.04.19. Includes latest changes. This version should be used until updated

Product name: Switch

Active Substance: Cyprodinil 37.5% w/w
Fludioxonil 25.0% w/w

Formulation: Water dispersible granule (WG)

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Authorisation Holder: Syngenta UK Ltd

Marketing Company: Syngenta Ireland Ltd

PCS No: 06414

User type/Function: Professional Use Only / Fungicide.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Indoor cucumber, gherkin, courgette	0.8 kg/ha in 1200l water/ha	3 per crop	2.4 kg/ha	7 days	3 days before harvest	<i>Mycosphaerella</i> , <i>Botrytis</i> Spp and <i>Fusarium</i>
Indoor tomato	1.0 kg/ha in 800l water/ha	3 per crop	3.0 kg/ha	10 days	3 days before harvest	<i>Botrytis</i>
Indoor pepper, aubergines	1.0 kg/ha in 600l water/ha	3 per crop	3.0 kg/ha	7 days	7 days before harvest	<i>Botrytis</i>
Indoor cress, endives, Lambs lettuce, lettuce, rocket, herbs, spinach, spinach beet & baby leaf brassica	0.6 kg/ha	2 per crop	1.2 kg/ha	10 days	7 days before harvest	<i>Botrytis</i>
Outdoor cress, endives, Lambs lettuce, lettuce, rocket, herbs, spinach, spinach beet & baby leaf brassica	0.8 kg/ha	2 per crop	1.6 kg/ha	10 days	7 days before harvest	<i>Botrytis</i>
Outdoor asparagus	1.0 kg/ha in 200l water/ha	3 per crop	3.0 kg/ha	7 days	After harvest	<i>Botrytis</i> Spp and <i>Stemphyllium</i>
Bulb onion, garlic, shallot	1.0 kg/ha in 200l water/ha	3 per crop	3.0 kg/ha	-	21 days before harvest	<i>Botrytis</i>

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Outdoor salad onion	1.0 kg/ha in 200l water/ha	3 per crop	3.0 kg/ha	7 days	14 days before harvest	<i>Botrytis</i>
Lupins	1.0 kg/ha	2 per crop	2.0 kg/ha	7 days	*	<i>Botrytis</i>
Beetroot, horseradish, parsnip, parsley root and salsify	0.8 kg/ha in 200l water/ha	3 per year	2.4 kg/ha	-	7 days before harvest	<i>Alternaria</i> and <i>Sclerotinia</i>
Outdoor celery	1.0 kg/ha	2 per crop	2.0 kg/ha	10 days	14 days before harvest	Leaf spot and <i>Sclerotinia</i>
Cherry, Plum	0.2 (during flowering) 0.6 (pre- harvest)	3 per year	1.8 kg/ha	-	7 days before harvest	Blossom wilt, fruit diseases
Indoor blueberry	0.8 kg/ha in 600l water /ha	3 per crop	2.4 kg/ha	10 days	3 days before harvest	<i>Botrytis</i> and <i>Anthraco</i> spp

* Straw from treated crops must not be used for animal feed or bedding

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application may only be made using a tractor mounted/trailer sprayer, handheld sprayer or broadcast air assisted sprayer.
- (4) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.
- (5) This approval extends to professional use only.
- (6) For environmental information refer to the product label but specifically:
To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

To protect aquatic organisms respect an unsprayed buffer zone of 10m to surface water bodies from broadcast air assisted sprayers. Aim spray away from water.

Aim spray away from water.

- (7) For operator protection refer to the product label.
- (8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Switch as a horticultural fungicide for the control of various diseases on the crops & situations listed above.

Resistance Management:

Use Switch as part of an Integrated Crop Management (ICM) strategy.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy.

Cyprodinil is classed: FRAC code 9.

Fludioxonil is classed FRAC code 12

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.