



Plant growth regulator for crop height reduction, lodging prevention and yield protection in cereals and grassland

An emulsifiable concentrate containing 250 g/L of trinexapac-ethyl (26.6 % w/w).



GROWTH REGULATOR

RISK AND SAFETY INFORMATION

Flammable liquid and vapour.
Causes serious eye irritation.
Harmful if inhaled.
May cause respiratory irritation.
May cause damage to organs through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.
Keep away from heat/sparks/open flames/hot surfaces. – No smoking.
Avoid breathing fumes/spray.
Wear protective gloves/ protective clothing/ eye protection/ face protection.
IF INHALED: remove victim to fresh air and keep at rest in a position comfortable for breathing.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.
Call a POISON CENTRE or doctor/physician if you feel unwell.
In case of fire: Use dry sand, dry chemical or alcohol-resistant foam to extinguish.
Collect spillage.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.



Repeated exposure may cause skin dryness or cracking.
Contains trinexapac-ethyl. May produce an allergic reaction.
To avoid risks to human health and the environment comply with the instructions for use.

FOR PROFESSIONAL USE ONLY | UFI: 3520-S0T1-K002-C3DP | PCS no. 04915

CONTENT: 1 litre



DISTRIBUTOR:
DHM Agrochemicals
Castle Lodge - Kilgobbin Road
Droghda D18R2W1
Telephone: (0)125 2237
www.dhm.ie

REGISTRATION HOLDER:
GLOBACHEM N.V.
Brusem Industriepark
Nichtwegberglaan 2019
B-3800 Sint-Truiden, Belgium
Tel: +32 11 78 57 17
Fax: +32 11 68 15 65
globachem@globachem.com
www.globachem.com



- ▶ An emulsifiable concentrate containing 250 g/L of trinexapac-ethyl (26.6 % w/w)
- ▶ PCS no. 04915



GROWTH REGULATOR

IMPORTANT INFORMATION
FOR USE ONLY AS AN AGRICULTURAL PLANT GROWTH REGULATOR

CROPS/SITUATIONS	MAXIMUM SINGLE AND TOTAL DOSE (L/ HA/CROP)	LATEST TIME OF APPLICATION
Winter wheat	0.4	Before flag leaf sheath extending stage (GS41)
Winter barley	0.6	Before flag leaf sheath extending stage (GS41)
Winter oats and spring oats	0.4	Before second node detectable stage (GS 32)
Grassland (seed crop)	0.8	Before second node detectable stage (GS 32)
Spring barley	0.5	Before third node detectable stage (GS 33)
Spring wheat, durum wheat, rye, triticale	0.4	Before third node detectable stage (GS 33)

READ THE LABEL BEFORE USE, USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE.
FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

BATCH NUMBER: SEE BOTTLE
READ ALL INSTRUCTIONS CAREFULLY BEFORE USE
PROTECT FROM FROST | STORE IN A COOL DRY PLACE
SHAKE WELL BEFORE USE

For 24 hour emergency information contact Globachem n.v.
Telephone: +32 11 78 57 17



Moxa® is a registered trademark for Globachem N.V.



5 425030 787816 >

ET100970_ED3-V1



IE



SAFETY PRECAUTIONS

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACE-SHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

AVOID ALL CONTACT WITH EYES.

WASH HANDS before eating and drinking/smoking/after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

ENVIRONMENTAL PROTECTION

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

STORAGE AND DISPOSAL

KEEP AWAY FROM FOOD DRINK AND ANIMAL FEEDING STUFFS

KEEP OUT OF REACH OF CHILDREN

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place/under lock and key/away from damp/sources of heat.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add the washings to the sprayer at the time of filling and dispose of safely.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

MOXA® is an emulsifiable concentrate formulation containing 250 g/L of imidacloprid. It is a growth regulator for crop height reduction, which can lead to lodging control and yield protection in all cereals and grassland. Treatment may lead to ears remaining erect through to harvest.

MIXING AND SPRAYING

Spray volume

Apply MOXA® in a minimum of 200 l/ha of water. Increased penetration will be obtained with an increase in water volume but the necessity for this will be dependent on crop growth stage and habit.

Spray nozzle

A MEDIUM spray quality is preferred for application of MOXA® (see BCPC guidelines). A spray pressure of 2–3 bar is recommended.

Mixing and spraying

Make sure the sprayer is set to give an even application at the correct volume. Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of MOXA® and continue agitation whilst adding the rest of the water. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. Agitate the mixture thoroughly before use and continue agitation during spraying. Take particular care to avoid overlapping of spray swaths. Wash out the sprayer thoroughly with water after use (at least 3 rinsings).

RESTRICTIONS

- Avoid spray drift onto neighbouring crops.
- Only use on crops at risk of lodging.
- Do not apply if rain or frost is expected or if the crop is wet.
- Do not spray crops under stress or to crops suffering from waterlogging, pest attack, disease or frost.

CROP SPECIFIC INFORMATION

Winter wheat

Apply 0.4 l/ha from the leaf sheath erect stage (GS30) to before the flag leaf sheath extending stage (GS41).

Winter barley

Apply 0.4 l/ha between leaf sheath erect stage and second node detectable stage (GS30–32).

OR

Apply 0.6 l/ha from the flag leaf just visible stage (GS37) to before the flag leaf sheath extending stage (GS41).

Spring barley

Apply 0.5 l/ha between leaf sheath erect stage (GS30) to before the third node detectable stage (GS33).

Spring wheat, Durum wheat, Rye, Triticale

Apply MOXA® at 0.4 l/ha between leaf sheath erect stage (GS30) to before the third node detectable stage (GS33).

Winter oats and spring oats

Apply MOXA® at 0.4 l/ha between the leaf sheath erect stage (GS30) to before the second node detectable stage (GS32).

Grassland (seed crops)

Apply MOXA® at 0.8 l/ha between the leaf sheath erect stage (GS30) to before the second node detectable stage (GS32).

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.