

# Basagran® SG

**A contact herbicide for post-emergence broad-leaved weed control in spring and winter field beans, broad, dwarf french, navy and runner beans, peas, linseed, narcissi and potatoes.**

A water soluble granule containing (87% w/w) bentazone as the sodium salt.

## Risk and Safety Information

### Danger

**Harmful if swallowed.**

**May cause an allergic skin reaction.**

**Causes serious eye damage.**

Wear protective gloves and eye face protection.  
Avoid breathing dust.

IF ON SKIN (or hair): Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTRE or doctor/physician.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

PCS No. : 05192



### Supplied by:

BASF Ireland Limited, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, UK, Telephone: 00 44 (0)161 475 3000

Emergency Information (24 hours freephone): 0049 180 227 3112

Technical Enquiries: 0044 (0)845 602 2553 (office hours)

### Authorisation holder:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport SK1 3GG, Telephone: 00 44 (0)161 475 3000

**3 kg**



**FOR USE ONLY AS AN AGRICULTURAL/ HORTICULTURAL HERBICIDE, as directed below:**

| Crops                                                                                | Maximum individual dose | Maximum total dose (per crop) | Latest time of application                                                         |
|--------------------------------------------------------------------------------------|-------------------------|-------------------------------|------------------------------------------------------------------------------------|
| Beans<br><i>Phaseolous</i> species<br>(Dwarf French beans, Runner beans, Navy beans) | 1.1 kg/ha               | 1.1 kg/ha                     | BBCH 17 (7 leaves unfolded)                                                        |
| Beans<br><i>Vicia faba</i> species<br>(Broadbeans and winter/spring beans)           | 1.1 kg/ha               | 1.1 kg/ha                     | BBCH 17 (7 leaves unfolded)                                                        |
| Peas<br>Combining and vining peas                                                    | 1.1 kg/ha               | 1.1 kg/ha                     | Before BBCH 39 (before flower buds can be found enclosed on the terminal shoot)    |
| Linseed                                                                              | 1.1 kg/ha               | 1.1 kg/ha                     | BBCH 29<br>Before enclosed bud visible in leaf axis (BBCH 31)                      |
| Potatoes                                                                             | 1.1 kg/ha               | 1.1 kg/ha                     | BBCH 19 (9 leaves of main stem unfolded) or before shoots exceed 15 cm in height # |
| Narcissi                                                                             | 1.1 kg/ha               | 1.1 kg/ha                     | No applicable*                                                                     |

\* This product must not be applied to ornamental plant production (narcissi) during flower bud formation.

# which ever comes first

**READ ALL PRECAUTIONS BEFORE USE****PCS No. : 05192****PRECAUTIONS**

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES, FACE PROTECTION (FACE SHIELD) AND SUITABLE RESPIRATORY PROTECTIVE EQUIPMENT when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WASH CONCENTRATE from skin or eyes immediately.

DO NOT BREATHE SPRAY.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Do not contaminate water with the product or its container. Do not clean application equipment near surface water / avoid contamination via drains from farmyards and roads.

**STORAGE**

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

Do not re-use container for any other purpose.

## COMPANY ADVISORY INFORMATION

This section provides additional advice on product use and is not part of the Product Label under the Plant Protection Product Regulation (EC) No 1107/2009.

### COMPANY ADVISORY INFORMATION

#### Water Stewardship – Bentazone

|               |                                                                                     | High Risk Areas*        | All Other Areas |
|---------------|-------------------------------------------------------------------------------------|-------------------------|-----------------|
| <b>Soils</b>  | Shallow soils (<30-35cm) and stony soils (>10% surface covered) on chalk/ limestone | Do not use              | Avoid use       |
|               | Shallow (<30-35cm) soils on sandstone                                               | Do not use              | Avoid use       |
|               | Soils with shallow groundwater (<1m below surface)                                  | Do not use              | Avoid use       |
|               | Soils with low organic carbon (<1%)                                                 | Do not use              | Avoid use       |
|               | Other soils                                                                         | OK to use               | OK to use       |
| <b>Timing</b> | Spring and Summer                                                                   | OK to use after 1 April | OK to use       |
|               | Autumn and Winter                                                                   | Do not use              | Do not use      |

\* Groundwater Source Protection Zones 1+2

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Basagran SG is a herbicide for post-emergence broad-leaved weed control in spring and winter field beans, broad, dwarf french, navy and runner beans, peas, linseed, narcissi and potatoes.

## Restrictions/Warnings

### Professional Use Only

The best weed control will be achieved if crops are sprayed when weeds are small and actively growing, provided the crop is within the correct growth stages, see below.

It is essential to achieve full cover of all weed surface areas, especially the growing point. Therefore adjust spray volumes and pressures according to the size and density of the weed populations and crop cover.

It is preferable to use the split dose if conditions are likely to be very warm, sunny or humid, if foliage is tender or if a sensitive variety is to be treated (see 'Mixing and Application' section).

Where weeds classed as moderately susceptible form an important part of the weed population, the split dose will only give acceptable control if the weeds are cotyledon only. If true leaves are present, a single application of 1 kg/ hectare should be made.

### Sequences/Tank mixes

Sequence with other spray chemicals:

Do not apply insecticides within 7 days of treatment with Basagran SG.

Basagran SG BEFORE a post emergence grass herbicide: leave an interval of at least 7 days between treatments.

Basagran SG AFTER a post emergence grass herbicide: leave an interval of 14 days and carry out a leaf wax test where relevant.

### Weather conditions

Do not apply Basagran SG to any crop that may have been subjected to stress conditions, such as herbicide or disease injury, excessive acidity/alkalinity, trace element deficiency, drought, waterlogging, widely fluctuating temperatures, or physical damage to the foliage as caused by abnormal wind, rain, hail or frost.

Do not spray under HOT SUNNY CONDITIONS when temperatures are at or above 21°C, particularly during the mid day period. When the above conditions occur, spraying should be delayed until the evening. This is particularly important when there has been a change to hot sunny weather following a cool cloudy period.

Do not spray if rain or frost is expected, nor if the foliage is wet. **A minimum period of 6 hours free from rain is required after application.**

Do not apply Basagran SG during periods of drought or unseasonably cold weather as unsatisfactory weed control may result.

## Crops

Check varietal tolerance of varieties before using.

A satisfactory leaf wax test using crystal violet marker dye, must be carried out before applying Basagran SG to peas.

DO NOT treat forage pea varieties or mange-tout.

Do not use on first early or seed crop potatoes.

Restrictions relating to certain varieties of potatoes should be strictly observed.

Check also all specific restrictions and warnings relevant to potatoes.

CONSULT PROCESSORS BEFORE USE.

## Application

Do not overlap spray swaths.

**Avoid spray drift onto neighbouring crops, particularly lettuce and sugar beet.**

The recommended water volume is 150-400 litres/hectare. The lowest volume of 150 litres/hectare can only be used when weeds are no larger than cotyledon stage, weed density is low to moderate and crop shading is negligible.

WASH EQUIPMENT thoroughly immediately after use. Fill the tank with clean water and leave overnight. Spray out before storage or using other products. Traces of the product may cause damage to susceptible crops sprayed later.

## Weed control

### Susceptibility rating

| Weed Name        | Basagran SG<br>1.1 kg/ha   |                                                 | Basagran SG<br>Split dose:<br>0.55 + 0.55 kg/ha |                                                 |
|------------------|----------------------------|-------------------------------------------------|-------------------------------------------------|-------------------------------------------------|
|                  | Weed susceptibility rating | Maximum susceptible growth stage of target weed | Weed susceptibility rating                      | Maximum susceptible growth stage of target weed |
| Amaranth, Common | MR                         | Up to 6 lvs                                     | MR                                              | Up to 2 lvs                                     |
| Black-bindweed   | MS                         | Up to 2 lvs                                     | -                                               | -                                               |
| Charlock         | S                          | Up to 6 lvs                                     | S                                               | Up to 2 lvs                                     |

| Weed Name          | Basagran SG<br>1.1 kg/ha   |                                                 | Basagran SG<br>Split dose:<br>0.55 + 0.55 kg/ha |                                                 |
|--------------------|----------------------------|-------------------------------------------------|-------------------------------------------------|-------------------------------------------------|
|                    | Weed susceptibility rating | Maximum susceptible growth stage of target weed | Weed susceptibility rating                      | Maximum susceptible growth stage of target weed |
| Chickweed, Common  | S                          | Up to 6 lvs                                     | S                                               | Up to 2 lvs                                     |
| Cleavers           | MS                         | Up to 6 lvs                                     | MS                                              | Cotyledons                                      |
| Dead-nettle, Red   | MR                         | Up to 6 lvs                                     | MR                                              | Cotyledons                                      |
| Fat-Hen            | MR                         | Up to 2 lvs                                     | MR                                              | Cotyledons                                      |
| Mayweed spp.       | S                          | Up to 6 lvs                                     | S                                               | Up to 2 lvs                                     |
| Nightshade, Black  | S                          | Up to 6 lvs                                     | S                                               | Up to 2 lvs                                     |
| Penny-cress, Field | S                          | Up to 6 lvs                                     | S                                               | Up to 2 lvs                                     |
| Persicaria, Pale   | S                          | Up to 6 lvs                                     | -                                               | -                                               |
| Radish, Wild       | MS                         | Up to 6 lvs                                     | -                                               | -                                               |
| Redshank           | S                          | Up to 4 lvs                                     | S                                               | Up to 2 lvs                                     |
| Shepherd's-purse   | S                          | Up to 6 lvs                                     | S                                               | Up to 2 lvs                                     |

S: Susceptible , MS: Moderately Susceptible, MR: Moderately Resistant

## Crop Specific Information

### Dwarf French, Navy and Runner Beans

#### Time of Application

Apply Basagran SG in runner, dwarf French any navy beans when the crop has between two and seven trifoliate leaves.

Crop selectivity is partly dependent on adequate leaf wax formation. When conditions which reduce leaf wax occur the application should be delayed by at least 5 to 7 days to await recovery.

Basagran SG may cause transient scorch, which can be prolonged with applications later than the three trifoliate leaf stage.

A range of varieties has been tested for tolerance. However, if unsure, growers should check a small area in the first instance or consult BASF for an up to date list of tolerant varieties.

### Method of Application

Basagran SG can be applied either by a conventional single dose or preferably by a split dose treatment.

### Spring and Winter Field Beans and Broad-Beans

Basagran SG can be used alone or following a pre-emergence herbicide providing the crop is not adversely affected. Crop selectivity is partly dependent on adequate leaf wax formation. Frost, wet conditions, physical damage, disease or previous sprays can all reduce leaf wax and when this occurs application should be delayed by at least 5 to 7 days to await recovery.

### Methods of Application

Basagran SG can be applied either by a conventional single dose or by a split dose treatment. The correct crop and weed growth stages are given in the following table. See 'Mixing and Application' section.

**The single dose** is usually the preferred treatment for the winter bean crop due to overwintered weeds being more advanced.

**The split dose** is the preferred method in spring field beans provided that the weed and crop stage are suitable.

### Rates and Timing – Field and Broad Beans

| Crop                                       | Application method       | Rate kg/hectare                  | Crop stage                                                                                                   | Weed stage                                     |
|--------------------------------------------|--------------------------|----------------------------------|--------------------------------------------------------------------------------------------------------------|------------------------------------------------|
| Spring and winter Field Beans, Broad Beans | Split dose <sup>1</sup>  | 0.55kg/ha followed by 0.55 kg/ha | 2 to 7 leaf pairs, max. height 15 cm                                                                         | cotyledon to 2 leaves only (optimum cotyledon) |
|                                            | Single dose <sup>2</sup> | 1.1kg/ha                         | 2 to 7 leaf pairs, max. height 15 cm<br><br>Broad Beans <sup>2</sup> optimum crop stage is 3 to 4 leaf pairs | cotyledon to 6 leaves only (optimum 2 leaves)  |

### Important notes:

<sup>1</sup> If conditions are conducive to increased herbicide activity and crop scorch, e.g. warm sunny conditions, tender foliage or more sensitive variety, then the 0.55/0.55 kg/hectare split should be used.

<sup>2</sup> Broad beans are more sensitive than field beans. The split dose treatment is always preferable to the single dose in this crop.

Processors must be consulted before use in broad beans.

### **Crop Effects – Field and Broad Beans**

Slight crop scorch can occur after spraying. This is seen as a blackening of leaf margins, particularly on older leaves – all subsequent new leaves being unaffected. A temporary vigour check may also occur under some conditions. These effects generally have no influence on yield. They can, however, be more pronounced when applications are made beyond the optimum growth stage, particularly when the single dose is used and/or adverse weather conditions prevail.

A range of varieties has been tested for tolerance. However if unsure, growers should check a small area in the first instance or consult EASF for an up to date list of tolerant varieties.

### Warnings

#### **Avoid overlapping spray bouts.**

Do NOT apply on crops earlier than the 2 leaf pair stage, or crops more than 15 cm tall or if flower buds are visible.

### **Peas**

Basagran SG may be used following pre-emergence broad-leaf herbicides provided that the crop is undamaged and showing no adverse symptoms of herbicide application.

Avoid application to crops which are under stress from physical damage or disease, or other factors.

A satisfactory leaf wax test using crystal violet marker dye must be performed before applying Basagran SG on peas, particularly with early drillings which tend to have less leaf wax.

Slight scorch of the leaf margins and/or hormonal twisting may occur shortly after spraying but the crop will soon recover; yield and maturity will not be adversely affected.

### **Time of Application**

Apply Basagran SG, from when the peas have two nodes (2 fully expanded leaves) until before the flower buds can be found enclosed in the terminal shoot (equivalent to BBCH 39). The optimum timing is when the majority of weeds have germinated, but are still at the seedling stage.



## Rate of Application

Single application: apply Basagran at 1.1 kg/ha

Split application: apply Basagran at 0.55 kg/ha followed by 0.55 kg/ha

## Varietal Tolerances

DO NOT treat the varieties Carouby de Mausanne, Conquest, Danielle, Dinos, Filby, Fonado, Printana, Sherbourne, Turon, Santa or Vedette.

DO NOT treat forage pea varieties or mange-tout.

A range of varieties has been tested for tolerance. However if unsure, growers should check a small area in the first instance or consult BASF for an up to date list of tolerant varieties.

## Linseed

Since linseed offers poor weed competition, weed control is important, especially in the early growth stages.

Basagran SG may be used alone or following pre-emergence herbicides, provided that the crop is showing no adverse symptoms.

## Time of Application

Apply when the crop is between 2 leaf stage and before the flower buds are visible (up to and including BBCH 29).

The optimum timing is when the majority of weeds have germinated and are still in the seedling stage. As a guide, this is usually when the crop is between 7½-20 cm tall.

## Methods of Application

Basagran SG can be applied either by a conventional single dose or by a split dose treatment.

## Narcissi

A weed control programme of a suitable pre-emergence herbicide followed by Basagran SG is recommended.

## Time of Application

Apply Basagran SG at or after flowering but not during flower bud formation. If required, Basagran SG may be applied at any time after crop emergence except during flower bud formation. For optimum control weeds should be at the cotyledon to seedling stage.

## Methods of Application

Basagran SG can be applied either by a conventional single dose or by a split dose treatment.

## Varietal Tolerances

Selectivity has been satisfactory over many varieties. However, because there is such a large range, growers are advised to check a small area in the first instance.

### Potatoes

Basagran SG is a post-emergence treatment for use in maincrop and second early potatoes. It may be used alone or following a pre emergence herbicide in a programme of weed control provided the crop is undamaged and showing no adverse symptoms of herbicide application.'

Some leaf-yellowing or slight scorch may occur under certain conditions (see 'Varietal Tolerances' and 'Factors Affecting Crop Tolerance'). These effects are transient, only being present on leaves exposed to the spray. All subsequent growth is unaffected and yields are not adversely affected, provided that applications are within the following guidelines.

Do not irrigate for at least 24 hours following application. In dry conditions irrigation prior to Basagran SG application can be beneficial to ensure that weeds are actively growing. However, foliage must be dry before application. Avoid overlapping spray bouts.

### Time of Application

Apply Basagran SG from second leaf on main stem up to 60 cm when the majority of weeds have emerged and are at the cotyledon to seedling stage, but before most of the crop has reached a height of 15 cm.

Application when the crop is taller than 15 cm is not recommended because foliage scorch can be increased leading to a possible check to the crop vigour and yields. Weeds can also be shielded from the spray by crop foliage with later applications.

### Methods of Application

Basagran SG can be applied either by a conventional single dose or preferably by a split dose treatment. See 'Mixing and Application' section.

### Varietal Tolerances

| Variety | Safe To Treat | Borderline* | Do Not Treat |
|---------|---------------|-------------|--------------|
| Ailsa # |               | ✓           |              |
| Ambo #  | ✓             |             |              |
| Anna #  |               | ✓           |              |

| Variety       | Safe To Treat | Borderline* | Do Not Treat |
|---------------|---------------|-------------|--------------|
| Atlantic      |               | ✓           |              |
| Ausonia       | ✓             |             |              |
| Balmoral #    |               | ✓           |              |
| Bintje        |               |             | ✓            |
| Brodick #     | ✓             |             |              |
| Cabaret #     |               |             | ✓            |
| Cara          | ✓             |             |              |
| Carlingford # |               |             |              |
| Charlotte #   |               |             | ✓            |
| Costella      | ✓             |             |              |
| Cultra #      | ✓             |             |              |
| Desiree       |               | ✓           |              |
| Erntestolz #  |               |             |              |
| Estima        |               | ✓           |              |
| Fianna        |               |             | ✓            |
| Fontane #     |               |             | ✓            |
| Harmony #     |               | ✓           |              |
| Hermes #      |               | ✓           |              |
| Innovator #   |               | ✓           |              |

| Variety         | Safe To Treat | Borderline* | Do Not Treat |
|-----------------|---------------|-------------|--------------|
| Kind Edward     |               | ✓           |              |
| Lady Rosetta    |               |             | ✓            |
| Marfona         | ✓             |             |              |
| Markies #       |               |             | ✓            |
| Maris Peer      | ✓             |             |              |
| Maris Piper     | ✓             |             |              |
| Melody #        |               | ✓           |              |
| Morag           |               |             | ✓            |
| Morene          |               |             | ✓            |
| Nadine          | ✓             |             |              |
| Navan #         | ✓             |             |              |
| Obelix #        | ✓             |             |              |
| Pentland Crown  | ✓             |             |              |
| Pentland Dell   | ✓             |             |              |
| Pentland Squire | ✓             |             |              |
| Picasso #       |               | ✓           |              |
| Premiere #      |               | ✓           |              |
| Record          | ✓             |             |              |
| Rode Pipo       |               |             | ✓            |

| Variety         | Safe To Treat | Borderline* | Do Not Treat |
|-----------------|---------------|-------------|--------------|
| Romano          | ✓             |             |              |
| Rooster #       |               | ✓           |              |
| Russett Burbank |               |             | ✓            |
| Sante           |               |             | ✓            |
| Saturna #       | ✓             |             |              |
| Saxon #         | ✓             |             |              |
| Shepody         |               |             | ✓            |
| Shula #         | ✓             |             |              |
| Stemster        |               |             | ✓            |
| Stroma          |               |             | ✓            |
| Symphonia #     |               | ✓           |              |
| Valor #         | ✓             |             |              |
| Vekaro          | ✓             |             |              |
| Vanessa         | ✓             |             |              |
| Wilja           | ✓             |             |              |

\*These varieties have shown more foliage scorch than others. Particular attention should be paid to the crop stage and factors affecting crop tolerance.

# These varieties are recommended on the basis of limited (2 years) data.

DO NOT treat seed crops or first early varieties.

#### Factors Affecting Crop Tolerance

Basagran SG should not be applied to potatoes under hot, sunny conditions when temperatures are at or above 21°C, particularly during the mid day period. When these conditions are encountered, spraying should be delayed until evening. It should be noted that in general, crop foliage is more sensitive when there is a sudden change

to hot, sunny weather following a cool, cloudy period. This sensitivity diminishes after approximately three days.

Crop tolerance will also be reduced if the crop is under stress, e.g. from such factors as physical damage (as from high wind), heavy rain or hail, virus diseases, blackleg, nematodes, Rhizoctonia, excessive soil alkalinity or acidity, or frost either shortly before or after treatment. Wet foliage is prone to scorch by Basagran SG.

## Mixing and Application

### Mixing

Half fill the spray tank with clean water and start the agitation. Pour in the required amount of Basagran SG. Add the remainder of the water and continue agitation until spraying is completed.

### Application

All applications should be made as a **FINE** spray, as defined by BCPC, unless the highest water volumes are used when **FINE** or **MEDIUM** sprays are permissible. It is essential to achieve full cover of all weed surface areas, especially the growing point. Therefore **ADJUST SPRAY VOLUMES AND PRESSURES ACCORDING TO THE SIZE AND DENSITY OF THE WEED POPULATIONS AND CROP COVER.**

Basagran SG can be applied as a single dose or as a split dose in all recommended crops. The split dose will control susceptible weeds from cotyledon up to 2 true leaves while the single dose is effective on weeds up to 4-6 true leaf stage. The split dose method has generally given better weed control and enhanced crop safety over the conventional single dose but if weeds are larger than 2 true leaves, it is necessary to use the single dose method.

### Timing of Split Dose

The optimum timing for the first application is when the first flush of weeds are at the cotyledon stage. The second follow up dose should be applied within 7 to 10 days of the first dose, depending on the control achieved by the initial dose or the appearance of the second weed flush. See individual sections below for crop growth stage limitations.

**A maximum of two applications may be made: 0.55 kg/hectare followed by 0.55 kg/ hectare.**

The 0.55 kg/0.55 kg/hectare split is preferable if conditions are likely to be very warm, sunny or humid, if foliage is tender or if a sensitive variety is to be treated.

The recommended water volume is 150-220 litres/hectare. The lowest volume of 150 litres/hectare can only be used when weeds are no larger than cotyledon stage, weed density is low to moderate and crop shielding is negligible.

## Single Dose

Where Basagran SG is recommended as a single application in this text, it should be applied at 1.1 kg/hectare.

When applied alone, the recommended water volume is 220-400 litres/hectare.

The lowest volume of 220 litres/hectare should only be used when all the following conditions apply:

- weeds at cotyledon to 2 leaves only
- weed density is low to moderate
- crop shielding is negligible

Flat fan or high pressure hollow cone nozzles are suitable under these conditions.

For other situations, 330-400 litres/hectare is recommended, preferably applied through flat fan nozzles only, operating at an optimum pressure of approximately 3-3.5 bar, in order to produce a fine penetrating spray.

## Compatibility

For details of compatibilities contact your distributor, local BASF representative or the BASF Technical Services Hotline: 0044 845 602 2553.

## Trade Mark Acknowledgments

Basagran is a BASF trademark.

## Additional Product Safety Information

This section does not form part of the product label under S.I. No. 159 of 2012.

The information on this label is based on the best available information including data from test results.

## Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



[agricentre.basf.ie/Basagran-SG-IE/MSDS](http://agricentre.basf.ie/Basagran-SG-IE/MSDS)

Alternatively, contact your supplier.

# Basagran<sup>®</sup> SG

**A contact herbicide for post-emergence broad-leaved weed control in spring and winter field beans, broad, dwarf french, navy and runner beans, peas, linseed, narcissi and potatoes.**

A water soluble granule containing (87% w/w) bentazone as the sodium salt.

## Risk and Safety Information

### Danger

**Harmful if swallowed.**

**May cause an allergic skin reaction.**

**Causes serious eye damage.**

Wear protective gloves and eye face protection.  
Avoid breathing dust.

IF ON SKIN (or hair): Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTRE or doctor/physician.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment, comply with the instructions for use.**

**PCS No. : 05192**



Supplied by:  
BASF Ireland Limited, 4th and 5th Floors, 2 Stockport Exchange, Railway Road,  
Stockport SK1 3GG, UK, Telephone: 00 44 (0)161 475 3000  
Emergency Information (24 hours freephone): 0049 180 227 3112  
Technical Enquiries: 0044 (0)845 602 2553 (office hours)

### Authorisation holder:

BASF plc, 4th and 5th Floors, 2 Stockport Exchange, Railway Road, Stockport  
SK1 3GG, Telephone: 00 44 (0)161 475 3000

**3 kg**

