



Terplex®

Plant growth regulator for use on cereals and grassland (seed crop)



An emulsifiable concentrate containing 200 g/L of trinexapac-ethyl (20.75 % w/w).

GROWTH REGULATOR



WARNING



RISK AND SAFETY INFORMATION

Harmful if swallowed
Causes serious eye damage
Toxic to aquatic life with long lasting effects

Keep out of reach of children.
Avoid breathing dust/fume/gas/mist/vapours/spray.

Wear protective gloves / eye protection.
IF IN EYES, Rinse cautiously with water for several minutes.
Remove contact lenses if present and easy to do, continue rinsing.
Immediately call a POISON CENTER or doctor/physician.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water / Avoid contamination via drains from farmyards and roads)

FOR PROFESSIONAL USE ONLY

PCS no. 05487

CONTENT: 5 lit. e



ET13953_ED1-V1

Registration holder:

Globechem NV
Brustem Industriepark
Lichtenberglaan 2019
B-3800 Sint-Truiden
Belgium
Tel: +32 11 78 57 17
Fax: +32 11 68 15 65
globechem@globechem.com
www.globechem.com



SAFETY PRECAUTIONS

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACE-SHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY with plenty of water and seek medical advice.

WASH HANDS before eating and drinking/smoking/after work.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

ENVIRONMENTAL PROTECTION

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

STORAGE AND DISPOSAL

KEEP AWAY FROM FOOD DRINK AND ANIMAL FEEDING STUFFS

KEEP OUT OF REACH OF CHILDREN

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place/under lock and key/away from damp/sources of heat.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add the washings to the sprayer at the time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

GENERAL INFORMATION

TERPLEX® is an emulsifiable concentrate formulation containing 200 g/L trinexapac-ethyl. It is a growth regulator for crop height reduction, which can lead to lodging control and yield protection in all varieties of winter and spring wheat, winter barley, rye, triticale, and grassland (seed crops).

Treatment may lead to ears remaining erect through to harvest.

CROP SPECIFIC INFORMATION

TERPLEX® can be used in the following crops at the dose rates mentioned below.

Winter barley: Apply 0.5 L/ha between the end of tillering and the second node detectable stage (BBCH 29-32).

Winter wheat: Apply 0.5 L/ha between the end of tillering and

Spring wheat:

the flag leaf fully emerged stage (BBCH 29-39). Apply 0.5 L/ha between the leaf sheath erect stage and the second node detectable stage (BBCH 30-32).

Rye:

Apply 0.5 L/ha between the six tillers detectable stage and the second node detectable stage (BBCH 26-32).

Triticale:

Apply 0.5 L/ha between the five tillers detectable stage and the second node detectable stage (BBCH 25-32).

Grassland (seed crops): Apply 0.6 L/ha between the first node detectable stage and the third node detectable (BBCH 31-33).

MIXING AND SPRAYING

Spray volume

Apply TERPLEX® in a minimum of 200 L/ha water. Increased penetration will be obtained with an increase in water volume but the necessity for this will be dependent on crop growth stage and habit.

Spray nozzles

A MEDIUM spray quality is preferred for application of TERPLEX® (see RCPC guidelines). A spray pressure of 2-3 bar is recommended.

Mixing and spraying

Make sure the sprayer is set to give an even application at the correct volume. Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of TERPLEX® and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying. Take particular care to avoid overlapping of spray swaths. Wash out the sprayer thoroughly with water after use (at least 3 rinsings).

RESTRICTIONS

Only use on crops at risk of lodging.

Avoid spray drift onto neighbouring crops.

Do not apply if rain or frost is expected or if the crop is wet.

Do not spray crops under stress or to crops suffering from waterlogging, pest attack, disease or frost.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.



- ▶ **Plant growth regulator for use on cereals and grassland (seed crop)**
- ▶ An emulsifiable concentrate containing 200 g/L of trinexapac-ethyl (20.75 % w/w).
- ▶ PCS no. 05487



GROWTH REGULATOR

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL PLANT GROWTH REGULATOR.

CROPS	MAXIMUM SINGLE DOSE	MAXIMUM NO. OF APPLICATIONS	MAXIMUM TOTAL DOSE	LATEST TIME OF APPLICATION
Winter wheat & Winter barley	0.5 L/ha	-	0.5 L/ha/crop	Flag leaf stage: flag leaf fully unrolled, ligule just visible (BBCH 39)
Grassland (seed crop)	0.6 L/ha	-	6 L/ha/year	Third node detectable Stage (GS 33)
Spring wheat, rye & triticale	0.5 L/ha	-	0.5 L/ha/crop	Third node detectable Stage (GS 33)


Other specific restrictions: grassland (seed crop) uses.

This product must not be used on grass seed crops that will be grazed by livestock or cut for fodder.

Treated grassland must not be grazed or cut for fodder.

READ THE LABEL BEFORE USE, USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

READ ALL INSTRUCTIONS CAREFULLY BEFORE USE | BATCH NUMBER: SEE BOTTLE
PROTECT FROM FROST | STORE IN A COOL DRY PLACE | SHAKE WELL BEFORE USE



FOR PROFESSIONAL USE ONLY

For 24 hour emergency information contact Globachem n.v. Telephone: +32 11 78 57 17



5 404020 112937 >

ETI3954_ED1-V1

