



Hygeia

AGROXONE 50 WEEDKILLER

A soluble concentrate containing 500 g/L (44.25% w/w) MCPA as the dimethylamine salt.
A selective herbicide for the control of many broad-leaved weeds in established grassland.

**PROTECT FROM FROST
FOR PROFESSIONAL USE ONLY**

**MAX DOSE
2.7L/HA**
READ LABEL FOR DETAILS

SELECTIVE HERBICIDE FOR BROADLEAVED WEEDS



DANGER

RISK AND SAFETY INFORMATION

Harmful if swallowed.

Causes serious eye damage.

Wash hands thoroughly after handling.

Do not eat, drink or smoke when using this product.

Wear protective gloves/ protective clothing/ eye protection/ face protection.

IF SWALLOWED: Call a POISON CENTER or

doctor/physician if you feel unwell. Rinse mouth.

IF IN EYES: Rinse cautiously with water for several

minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Dispose of contents/ container to a licensed hazardous-waste disposal contractor or collection site except for empty triple rinsed containers which can be disposed of as non hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use

PCS No 05500

ADDITIONAL SAFETY PHRASES: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies. Direct spray away from water. Extreme care must be taken to avoid spray drift onto non-crop plants outside the target area. Livestock must be kept out of treated areas until poisonous weeds such as ragwort have died and become unpalatable. Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).
DO NOT Apply if rain is forecast within 48 hours. DO NOT Apply to saturated soils.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE:

Crop	Max individual dose	Max Total Dose	Latest timing
Established Grassland	2.7 L/ha	5.4 L product/ha/year	N/A
Grass seed crop	2.7 L/ha	2.7 L product/ha/year	5 weeks before heading

Other specific restriction

Do not apply by hand-held equipment. This product must not be applied between the end of September and beginning of March.

Do not apply in volumes less than 200 litres of water per hectare.

10 Litre e



5 390182 044035 >

MARKETED BY:

Hygeia Chemicals Ltd.

Oranmore, Co. Galway, Ireland.

Tel: 091-794722 Fax: 091-794738

Email: enquiries@hygeia.ie Web: www.hygeia.ie

Irish 24 hour emergency number:

National Poisons Centre in Ireland: 01-8092566

AUTHORISATION HOLDER:

Nufarm UK Limited, Wyke Lane, Wyke, Bradford,

West Yorkshire, BD12 9EJ, UK

DIRECTIONS FOR USE

RESTRICTIONS

- DO NOT roll or harrow within a week of spraying.
- DO NOT mow within four days of application.
- DO NOT treat newly established grass less than one year old.
- DO NOT treat grass suffering from drought, disease, during freezing conditions when rain is imminent or other adverse factors.
- Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area

- DO NOT spray in windy weather, avoid drift on to neighbouring crops, especially beans, beet, brassicas, carrots, legumes, lettuce and tomatoes which are very susceptible.
- WASH EQUIPMENT thoroughly immediately after use. Rinse with water three times, drain and allow to dry. Traces of herbicide left in the sprayer may damage susceptible crops if these are subsequently sprayed using the same equipment.
- Do not apply in volumes less than 200 litres of water per hectare

CROP SPECIFIC INFORMATION

Always spray when the crop is actively growing.

ESTABLISHED GRASSLAND Application rate: 2.7 L product/ha
 Maximum total dose: 5.4 L product/ha

Use on established grassland and leys not less than 1 year old. Apply the product in spring and summer when growing conditions are favorable. Spray grass for hay or silage 3-4 weeks before cutting. Clover may be checked but recovers by the following Spring. Follow-up applications may also be needed where new seedling weeds appear. An interval of 4-6 weeks should elapse between successive applications if weeds persist.

A top dressing ten days before treatment is recommended to assist kill of weeds and subsequent recovery of the sward. Annual weed species will be best controlled if spraying is done while the majority of weeds are seedlings. Perennial weeds should be sprayed during their period of maximum growth, usually when the flower buds are beginning to form. The response of perennial weeds to treatments are often variable with only the aerial parts killed, though often suppression will occur. The recovery of the weeds will be reduced if the grass is growing vigorously at the time of treatment. Apply in 200 - 400 L water/ha.

GRASS SEED CROP Application rate:
 2.7 L product/ha Maximum total dose: 2.7 L product/ha

The safe period for treatment occurs when the grasses have at least four leaves and have begun to tiller but at least five weeks preceding ear emergence. The best time varies according to the species and strain of grass concerned. Do not apply more than 2.7 L/ha of the product. Apply in 200 - 400 L water/ha.

MIXING AND SPRAYING

Half-fill the tank with clean water and add the required quantity of the product, and add the remainder of the clean water with gentle agitation which should continue until the contents are thoroughly mixed. Triple rinse containers with water and add washing to the spray tank.

TANK CLEANING

WASH EQUIPMENT thoroughly immediately after use. Rinse with water three times, drain and allow to dry. Traces of herbicide left in the sprayer may damage susceptible crops if these are subsequently sprayed using the same equipment.

COMPATIBILITY

The product can be tank-mixed with other pesticides, please consult your distributor.

RESISTANCE MANAGEMENT

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered to be resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures.

SUSCEPTIBILITY	WEED SPECIES	TIMING
Susceptible (Consistently good control of both roots and shoots)	Greater Plantain	
	Ribwort Plantain	
Moderately Susceptible (Weeds suppressed and top-growth usually killed)	Buttercup, Meadow	Treat in spring or early summer
	Cat's Ear	
	Cress, Hoary	
	Daisy, Common	
	Hawkbit, Autumn	
	Knapsweed, Common	
	Ragwort, Common	Spray in spring when in rosette stage before flower spike starts to grow
	Rush, Compact	Spray in April to June. Cut and remove stems either four weeks before or after treatment.
	Rush, Soft	Treat as Compact Rush
	Self Heal	
Moderately Resistant (Top growth only may be killed)	Thistle, Creeping	Treat at early flower bud stage
	Thistle, Spear	
	Buttercup, Bulbous	Treat in autumn on new leaf or in spring
	Buttercup, Creeping	Treat in spring or early summer
	Colt's Foot	
	Dandelion	
	Meadowsweet	
	Rush, Hard	Treat as Compact Rush
	Sorrel, Common	
	Sorrel, Sheep's	
	Thistle, Dwarf	
	Yarrow	

TERMS AND CONDITIONS OF SUPPLY, SALE OR USE All goods supplied by us are high grade and we believe them to be suitable for the purpose for which we expressly supply them: but as we cannot exercise any control over their mixing, use or application which may affect the performance of the goods all conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us or our Associate Companies for any damage or injury whatsoever arising from their storage, handling, re-application or use. These conditions cannot be varied by our staff, our agents or the re-sellers of the product whether or not they supervise or assist in the use of such goods.