


MAVITA[®] 250 EC

PCS 05941

A fungicide that controls a wide range of diseases of winter and spring oilseed rape, Brussels sprouts, head cabbage, kale, collard, broccoli, calabrese and cauliflower.

An emulsifiable concentrate containing 250 g/l (23.4% w/w) difenoconazole. Also contains solvent naphtha.



SAFETY INFORMATION

Danger
May be fatal if swallowed and enters airways.
Causes serious eye irritation.
Very toxic to aquatic life with long lasting effects.
Keep out of reach of children.
Avoid release to the environment.
Wear protective gloves/protective clothing/eye protection/face protection.
IF SWALLOWED: immediately call a POISON CENTRE or doctor/physician.
IF IN EYES: rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Do NOT induce vomiting. Collect spillage. Dispose of contents/ container to a licensed hazardous waste disposal contractor or collection site except for empty, triple rinsed, clean containers which can be disposed of as non-hazardous waste. To protect aquatic organisms, respect an unsprayed buffer zone of 5 m to surface waters.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS 05941

XXXXXXXXXX

FR R3.0 JAN18

5 litres e

To access the Safety Data Sheet for this product scan the QR code or use the weblink: <http://XXXXXXXXXX>
- Alternatively, contact your supplier.



Scan for
Safety Data
Sheet

ADAMA

MAVITA® 250 EC**PCS 05941****PROTECT FROM FROST
SHAKE WELL BEFORE USE****DIRECTIONS FOR USE:**

see leaflet for full instructions.

Adama Agricultural Solutions UK Ltd

Unit 15, Thatcham Business Village, Colthrop Way, Thatcham, Berkshire RG19 4LW

Telephone: +44 (0)1635 860555 | **Technical Helpline:** +44 (0)1635 876622**www.adama.com | ukenquiries@adama.com****For advice on medical emergencies, fires or major spills telephone
the UK National Chemical Emergency Centre on +44 (0)1865 407333**

Batch No.: see packaging

ADDITIONAL SAFETY PRECAUTIONS**Operator Protection**

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces during spray application.

WASH SPLASHES from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

FOR USE VIA TRACTOR MOUNTED/TRAILED SPRAYERS ONLY.

Environmental Protection

To protect aquatic organisms, respect an unsprayed buffer zone of 5 m to surface waters.

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

Storage and Disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

CONDITIONS OF USE
FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose	Maximum number of treatments	Maximum total dose	Latest time of application
Oilseed rape	0.5 L/ha	-	1.0 L/ha	Before end of flowering (GS 69)
Brussels sprouts, cabbage, broccoli/calabrese, cauliflower, kale and collard	0.5 L/ha	-	1.0 L/ha	21 days before harvest

Other specific restrictions:

A minimum interval of 14 days must be observed between applications to brassica crops.

Applications to brassica must only be made to developed canopy and not before growth stage BBCH 41 as described below:

- Cabbage: heads begin to form, the 2 youngest leaves do not unfold.
- Kale, collard: 10% of the leaf mass typical for the variety reached.
- Brussels sprouts: lateral buds begin to develop.
- Cauliflower, broccoli, calabrese: heads begin to form, width of growing tip >1cm.

This leaflet/booklet is part of the approved label.

For advice on medical emergencies, fires or major spills telephone
the UK National Chemical Emergency Centre on +44 (0)1865 407333

**A fungicide that controls a wide range of diseases of winter and spring oilseed rape,
Brussels sprouts, head cabbage, kale, collard, broccoli, calabrese and cauliflower.**

An emulsifiable concentrate containing 250 g/l (23.4% w/w) difenoconazole.

Also contains solvent naphtha.

SAFETY INFORMATION

Danger

May be fatal if swallowed and enters airways.

Causes serious eye irritation.

Very toxic to aquatic life with long lasting effects.

Keep out of reach of children.

Avoid release to the environment.

Wear protective gloves/protective clothing/eye protection/
face protection.

IF SWALLOWED: immediately call a POISON CENTRE or
doctor/physician.

IF IN EYES: rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue
rinsing.

Do NOT induce vomiting.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste
disposal contractor or collection site except for empty, triple
rinsed, clean containers which can be disposed of as
non-hazardous waste.

To protect aquatic organisms, respect an unsprayed buffer
zone of 5 m to surface waters.

**To avoid risks to human health and the environment,
comply with the instructions for use.**

PCS 05941

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

An emulsifiable concentrate containing 250 g/l (23.4% w/w) difenoconazole. MAVITA® 250 EC is a DMI fungicide with contact and systemic activity against a wide range of diseases.

RESISTANCE MANAGEMENT

MAVITA 250 EC is a DMI fungicide. Strains of light leaf spot resistant to DMI fungicides are known to exist. To avoid development of resistance, apply product protectively in response to disease forecasts. Where possible, when light leaf spot is present, avoid the use of DMI based fungicides when targeting other diseases such as *Sclerotinia* at mid flowering.

DISEASES CONTROLLED

On oilseed rape, MAVITA 250 EC controls light leaf spot, leaf and pod spot and stem canker.

On Brussels sprouts, cabbage, broccoli, calabrese, kale, collard and cauliflower, MAVITA 250 EC controls leaf spot and ring spot.

CROP SPECIFIC INFORMATION

Winter and spring sown oilseed rape

MAVITA 250 EC controls light leaf spot, leaf and pod spot (*Alternaria* spp) and stem canker (*Phoma lingam*).

Light leaf spot

Spray protectively with 0.5 litres MAVITA 250 EC per ha from the 4 expanded true leaf stage in the autumn where there is a high risk of disease. A repeat spray may be made in the spring at the beginning of stem extension if visual symptoms develop.

In areas of low disease risk, spraying may be delayed until the spring when crops should be sprayed at the first sign of disease.

For moderate control of light leaf spot only, a 2 spray programme of 0.25 litres per ha may be used, starting in the autumn.

Leaf and pod spot (*Alternaria* spp)

For effective control spray 0.5 litres MAVITA 250 EC per ha at the end of flowering.

Stem canker (*Phoma lingam*)

Spray 0.5 litres MAVITA 250 EC per ha in the autumn after the 4 expanded true leaf stage when disease symptoms first occur. Repeat with an additional application of 0.5 litres per ha if foliar disease symptoms redevelop in late winter or early spring with a maximum total dose of no more than 1.0 litre per ha.

ALTERNATIVELY

A 2 spray programme of 0.25 litres MAVITA 250 EC per ha may be used, starting in the autumn when disease symptoms first occur, with the second application 4-6 weeks after the initial application. Repeat with an additional application of 0.5 litres per ha if foliar disease symptoms redevelop in late winter or early spring with a maximum total dose of no more than 1.0 litre per ha.

Brussels sprouts, cabbage, broccoli, calabrese, kale, collard and cauliflower

MAVITA 250 EC controls leaf spot (*Alternaria* spp) and ring spot. A 2 spray programme should be used commencing at the first sign of disease but not before the canopy has developed to growth stage BBCH 41 as described below.

- Cabbage: heads begin to form, the 2 youngest leaves do not unfold
- Kale, collard: 10% of the leaf mass typical for the variety reached
- Brussels sprouts: lateral buds begin to develop
- Cauliflower, broccoli, calabrese: heads begin to form, width of growing tip >1cm

A minimum interval of 14 days must be observed between applications. Best control will result from spraying before disease is established in the crop.

MIXING AND SPRAYING

Mixing

Make sure the sprayer is set to give an even application at the correct volume.

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of MAVITA 250 EC and continue agitation while adding the rest of the water.

Agitate the mixture thoroughly before use and continue agitation during spraying.

Thoroughly wash all spray equipment with water immediately after use.

Spray Volume

Oilseed rape: 200 litres water per hectare

Brassica: 400 litres water per hectare

Spraying

Apply as a MEDIUM quality spray as defined. A spray pressure of 2-3 bar is preferred.

DISCLAIMER/CONDITIONS OF SUPPLY

The specified properties of our products and the mode of application stated on this label have been established on the basis of research and experience. Products conform to specification at the time of delivery but, as we exercise no control over their subsequent storage, handling, mixing or use or the weather conditions before, during and after application, all of which may affect the performance of the products, no responsibility or liability will be accepted by us or our re-sellers for any failure in performance, damage or injury to person or property whatsoever arising from the storage, handling, application or use of the products. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in or make recommendations concerning the use of such products. We recommend you contact your dealer to request advice on the suitability of this product for any new and/or unusual growing methods or for new varieties not listed on this label.

Marketed by

Adama Agricultural Solutions UK Ltd
Unit 15, Thatcham Business Village
Colthrop Way, Thatcham, Berkshire RG19 4LW
Telephone: 00 44 1635 860555
Technical Helpline: 00 44 1635 876622
www.adama.com
Email: ukenquiries@adama.com

Authorisation Holder

Syngenta UK Limited
CPC 4, Capital Park, Fulbourn, Cambridge CB21 5XE

MAVITA® is a registered trademark of a company of the Adama Group.