

For the control of annual grass weeds and annual broad-leaved weeds in
Winter barley, Winter wheat and Potatoes (seed and ware).

Contains 800 grams per litre
prosulfocarb as an emulsifiable
concentrate formulation.

**FOR PROFESSIONAL USE
ONLY**

Read label before use

Keep out of reach of children

**SHAKE WELL BEFORE USE
PROTECT FROM FROST**

Batch No. and manufacturing date:
see packing

10 Litres e

Danger

May be fatal if swallowed and enters airways.

Causes skin irritation.

May cause an allergic skin reaction.

Causes serious eye irritation.

Very toxic to aquatic life with long lasting effects.

Avoid breathing dust/ fume/ gas/ mist/ vapours/ spray.

Wash hands thoroughly after use

Wear protective gloves/ eye protection/ face
protection.

IF SWALLOWED: Immediately call a POISON
CENTRE or doctor/physician.

Do NOT induce vomiting.

IF ON SKIN: Wash with plenty of soap
and water

If skin irritation or rash occurs: Get
medical advice/ attention.

IF IN EYES: Rinse cautiously with
water for several minutes. Remove
contact lenses, if present and easy to do.

Continue rinsing

If eye irritation persists: Get medical
advice/attention

Dispose of contents/ container to a licensed hazardous-
waste disposal contractor

or collection site except for empty triple rinsed clean
containers which can be

disposed of as non-hazardous waste.

Do not contaminate water with the product or its container (Do not
clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads)

**In the event of emergency, call the National Poisons Information Centre,
Beaumont Hospital at 01 809 2166 or 01 837 9964**

If medical advice is needed, have product container or label at hand.

**To avoid risks to human health and the environment,
comply with the instructions for use.**

PCS 06482

Manufacturer: Barclay Chemicals Manufacturing Ltd.,

Damastown Way, Damastown Industrial Park, Mulhuddart, Dublin 15, Ireland.

Tel: +353 1 8112900 Fax: +353 1 8224678 E-mail: info@barclay.ie Website: www.barclay.ie

Approval Holder: Barclay Chemicals (R&D) Ltd. Contact details as above.

Copyright © Barclay Chemicals (R&D) Limited, 2019.

© Crozier is a registered trademark of Barclay Chemicals (R&D) Ltd

To protect aquatic organisms, respect an unsprayed buffer zone of 5m to surface water bodies.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Take off contaminated clothing and wash before reuse

Contaminated work clothing should not be allowed out of the workplace

Wash contaminated clothing before reuse

Store locked up

Collect spillage

IMPORTANT INFORMATION			
FOR USE ONLY AS AN AGRICULTURAL HERBICIDE			
Crops	Maximum individual dose (L product/ha)	Maximum number of treatments per crop	Latest time of application
Wheat (winter), barley (winter)	5 litres (pre-emergence) OR	One per crop	Pre-emergence
	3 litres (up to GS22)		Early tillering (GS 22)
Potatoes (seed and ware)	5 litres	One per crop	At emergence (soils rising over emerging potato shoots)

Other specific restrictions:
Do not apply by hand-held equipment.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTISE FOR PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: this information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

RESTRICTIONS

Do not apply to crops under stress or to crops suffering from waterlogging, pest attack, disease, frost or the effects of high diurnal temperature changes. Transient yellowing can occur although crops fully recover.

For cereals, seed must be covered by 3cm of soil and for best results apply to a firm, moist seedbed free of clods.

DO NOT sow field beans or broad beans within 12 months of application.

WEEDS CONTROLLED

GRASSWEEDS CONTROLLED	
Susceptible	
Rough stalked meadow grass	Pre-emergence
Moderately susceptible	
Annual meadow grass	Up to 3 true leaves
Loose silky bent	Pre-emergence

CROZIER used pre-emergence will reduce blackgrass and Italian ryegrass populations. It should only be used as part of an appropriate management strategy involving sequences with products of alternative modes of action and the use of cultural techniques.

BROAD-LEAVED WEEDS CONTROLLED	
Susceptible	
Ivy leaved speedwell	Susceptible at up to cotyledon stage
Black nightshade	Pre-emergence
Chickweed	Susceptible at up to 2 true leaves
Forget Me Not	Pre-emergence
Cranesbill	Pre-emergence
Red dead nettle	Pre-emergence
Common field speedwell	Susceptible at up to cotyledon stage
Green field speedwell	Susceptible at up to cotyledon stage
Wall speedwell	Pre-emergence
Moderately susceptible	
Cleavers	Moderately susceptible at up to 1 whorl
Resistant	
Field pansy	Resistant at emergence

Off label approvals				
Crop	Max. Ind. dose	Max. no. of applications	Max. total dose	Latest time of application
Field onion, garlic, shallot, leek	5.0 l/ha	-	5.0 l/ha	Before 5th leaf is visible
Salad onions, chives	5.0 l/ha	-	5.0 l/ha	Before 3rd true leaf
Field celery	5.0 l/ha	1 per crop	5.0 l/ha	Growth stage BBCH 11. 50 days before harvest
Carrot, parsnip, parsley root, salsify, horseradish, beetroot	5.0 l/ha	-	5.0 l/ha	Growth stage BBCH 11. 80 days before harvest
Field Bean	5.0 l/ha	1	5.0 l/ha	Pre-emergence
Triticale, Rye	3.0 l/ha	1	3.0 l/ha	Before early tillering (GS 22)
Outdoor woody ornamentals & forest nursery	5.0 l/ha	-	5.0 l/ha	30th November to 1st March application only (During dormancy)

Weed Resistance

Strains of some annual grasses (e.g. black-grass, wild-oats, and Italian rye-grass) have developed resistance to a range of herbicides which may lead to poor control from one or more product or mode of action.

A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

CROZIER should only be used for control of herbicide resistant strains of annual grasses as part of an appropriate management strategy, including sequences with herbicides of alternative modes of action and the use of cultural techniques.

CROP SPECIFIC INFORMATION

Winter Cereals

CROZIER applied pre-emergence may infrequently slow crop emergence. This effect is transient and has been demonstrated not to adversely affect yield.

Potatoes

For control of a wider spectrum of weeds use CROZIER in mixture with approved formulations of products containing the active ingredient metribuzin, only as a pre-emergence application. Please consult company literature for specific product and best use guidelines. Always observe full label restrictions for any tank mix partner.

Timing

Winter Cereals

CROZIER can either be used at pre-emergence of the crop (5 l/ha) or at post-emergence of the crop up to GS 22 (3 l/ha). One application per crop.

Early and Maincrop Potatoes

CROZIER may be applied pre-emergence or at emergence of the crop (soil rising over emerging potato shoots). Complete ridge formation before application of CROZIER and do not disturb treated soil after application.

Rates of Use

Winter Cereals pre-emergence

5 litres of product per hectare. One application per crop.

OR

Winter Cereals early post-emergence up to GS22

3 litres of product per hectare. One application per crop.

Early and Maincrop Potatoes up to GS11

5 litres of product per hectare. One application per crop.

FOLLOWING CROPS

Winter Cereals, Early and Maincrop Potatoes

Do not sow field beans or broad beans within 12 months of application.

In the case of winter cereal crop failure, Winter Wheat or Winter Barley may be sown immediately in the autumn.

The following crops may be sown in the spring after winter cereal crop failure

Without ploughing

Sunflowers

Maize

Flax

Spring oats

Spring barley

Spring wheat

Spring peas

Spring oilseed rape

Soya beans

With ploughing

Carrots
Lettuce
Onions
Sugar beets
Potatoes

Do not sow

Field beans
Broad beans

The following crops may be sown in the autumn after potato crop failure or normal harvest:

Without ploughing

Winter oats
Winter barley
Winter wheat

MIXING AND SPRAYING**MIXING**

Fill the spray tank with half the required volume of clean water and start agitation.

Add the required amount of CROZIER and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying.

SPRAYING

Apply CROZIER in a water volume of 200-400 litres per hectare.

Apply CROZIER using a conventional fan nozzle producing a medium spray quality. A spray pressure of 2.0-3.0 bars is recommended.

CROZIER is rainfast after 1 hour.

Method of application: Tractor mounted / trailed sprayer

WASHING OUT PROCEDURE

Immediately after use, clean the spray equipment thoroughly. Drain the system completely and rinse spray tank, boom and nozzles two to three times with clean water until the foam and all traces of product have been removed.