

TRUSS

An emulsifiable concentrate containing 250g/l (25.5% w/w) trinexapac-ethyl per litre.

FOR USE ONLY AS AN AGRICULTURAL GROWTH REGULATOR

A growth regulator for winter and spring wheat, winter and spring barley, winter and spring oats, rye, triticale and durum wheat



FOR PROFESSIONAL USE ONLY
CLAYTON TRUSS: An emulsifiable concentrate formulation containing 250 g/l trinexapac-ethyl.

Signal Word: WARNING

Hazard Statements:

May cause an allergic skin reaction
Toxic to aquatic life with long-lasting effects.

Precautionary Statements:

Keep out of reach of children.
 Avoid breathing dust/fume/gas/mist/vapours/spray.
 Wear protective gloves/protective clothing/eye protection/face protection.
 IF ON SKIN: Wash with plenty of soap and water.
 If skin irritation or rash occurs, get medical advice.
 Collect spillage.
 Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste

Supplemental Information: Repeated exposure may cause skin dryness or cracking.

To avoid risks to human health and the environment comply with the instructions for use.

PCS 06494

In the event of emergency, call the National Poisons Information Centre, Beaumont Hospital at 01 809 2166 or 01 837 9964

Net Contents:

1 Litre e

SHAKE THOROUGHLY BEFORE USE
PROTECT FROM FROST

IMPORTANT INFORMATION: FOR USE ONLY AS AN AGRICULTURAL PLANT GROWTH REGULATOR

Crop	Maximum Individual Dose (product per hectare)	Maximum Total Dose (product per hectare per crop)	Latest time of application
Winter Wheat	0.4	0.4	Before flag leaf sheath extending stage (GS 41)
Spring Wheat	0.4	0.4	Before third node detectable stage (GS 33)
Winter Barley	0.6	0.6	Before flag leaf sheath extending stage (GS 41)
Spring Barley	0.5	0.5	Before third node detectable stage (GS 33)
Winter and Spring Oats	0.4	0.4	Before second node detectable stage (GS 32)
Rye, Triticale Durum Wheat	0.4	0.4	Before third node detectable stage (GS 33)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

SAFETY PRECAUTIONS

Operator protection

AVOID CONTACT WITH SKIN AND EYES.
 WEAR EYE/FACE PROTECTION when handling the concentrate.
 FOR USE BY TRACTOR MOUNTED/RALLED SPRAYERS ONLY

Environmental protection

Do not contaminate water with the product or its container.
 Do not clean application equipment near surface water.
 Avoid contamination of drains from farm yards and roads.

Storage and disposal

KEEP IN ORIGINAL CONTAINER, tightly closed in a safe place.
 RINSE THE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.
 Do not re-use container for any other purpose and dispose of safely.

Restrictions

Apply **Clayton Truss** only to healthy, actively growing crops.
 Do not apply during periods of frosty weather or when frost is imminent.
 Do not apply **Clayton Truss** to crops that are stressed by severe weather conditions, drought, frost, disease, insect damage, nutritional deficiency, etc.
 Do not apply if rain is expected or if the crop is wet.
 Avoid spray drift on to neighbouring crops.

CONDITIONS OF SUPPLY

All goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

GENERAL INFORMATION

CLAYTON TRUSS is a growth regulator for crop height reduction, which can lead to lodging control and yield protection in all varieties of winter and spring wheat, winter and spring barley, winter and spring oats, rye, triticale and durum wheat.

Treatment may lead to ears remaining erect through to harvest.

Mixing and Spraying

Make sure the sprayer is set to give an even application at the correct volume.
 Fill the spray tank with half the required volume of clean water and start agitation.
 Add the required amount of **CLAYTON TRUSS** and continue agitation whilst adding the rest of the water.
 Agitate the mixture thoroughly before use and continue agitation during spraying.
 Take particular care to avoid overlapping of spray swaths.
 Thoroughly wash all spray and measuring equipment with water and a wetting agent immediately after use.

MIXING AND SPRAYING

Spray Volume: Apply **CLAYTON TRUSS** in a recommended 200 l/ha of water. Increased penetration will be obtained with an increase in water volume but the necessity for this will be dependent on crop growth stage and habit.

Spray Nozzles: A medium spray quality is preferred for application of **CLAYTON TRUSS**. A spray pressure of 2-3 bar is recommended.

Spraying: Take particular care to avoid overlapping of spray swaths. Apply only using a ground sprayer.

CROPS

Winter Wheat:

Apply at 0.4 l/ha between the leaf sheath erect stage (GS30) but before the flag leaf extended stage (GS41).

Spring Wheat :

Apply at 0.4 l/ha from the leaf sheath erect stage (GS 30) but before the third node detectable stage (GS 33).

Winter Barley:

Apply at 0.4 l/ha between leaf sheath erect stage (GS30) but before the third node detectable stage (GS 33).

Or :-

Apply at 0.6 l/ha from the flag leaf just visible stage (GS 37) but before the flag leaf extending stage (GS 41).

Spring Barley:

Apply at 0.5 l/ha from the leaf sheath erect stage (GS 30) but before the third node detectable stage (GS 33).

Winter and Spring Oats:

Apply at 0.4 l/ha from the leaf sheath erect stage (GS30) but before the 2nd node detectable stage(GS32).

Rye, Triticale and Durum Wheat:

Apply at 0.4 l/ha between the leaf sheath erect stage (GS30) but before the 3rd node detectable stage (GS 33).

CROP FAILURE

In the event of crop failure for any reason, cereals and oilseed rape can be planted in soil treated with MODDUS 250 EC. Due to reduced activity via the root system and to its rapid degradation in soil, no problems with following crops are foreseen for this product.