

170,00 mm



DHM Agrochemicals

# KINGMAN

A fungicide for protection against leaf blotch (*Zymoseptoria tritici*) on wheat and triticale and leaf blotch (*Rhynchosporium secalis*) on barley and triticale.

A suspension concentrate formulation containing 500 g/l (39.5% w/w) folpet.  
Contains folpet and hexamethyltetramine.



## FOR PROFESSIONAL USE ONLY

WARNING UFI: M300-D039-6005-G0TD

May cause an allergic skin reaction.  
Suspected of causing cancer.  
Very toxic to aquatic life with long lasting effects.

Keep out of reach of children.  
Obtain special instructions before use.  
Keep only in original packaging.  
Wear protective gloves/protective clothing.  
IF ON SKIN: wash with plenty of soap and water.  
If exposed or feel unwell:

Call a POISON CENTRE or doctor, physician.  
Dispose of content, container to a licenced hazardous waste disposal contractor or collection site except for triple rinsed, empty containers which can be disposed of as non-hazardous waste.

To protect aquatic organisms respect an unsprayed buffer of 5 metres to surface water bodies.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No: 06666

**PCS No:  
06666**

Approval Holder and Distributor:  
**DHM Agrochemicals Ltd**  
Castle Lodge  
Kilgobbin Road  
Dublin 18  
Emergency Telephone  
number: 01 2952377.  
web: [www.dhm.ie](http://www.dhm.ie)

## SAFETY PRECAUTIONS

### OPERATOR PROTECTION

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES

when handling the concentrate or handling contaminated surfaces.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating or drinking and after work. IF YOU FEEL UNWELL, seek medical advice immediately (show label where possible).

### ENVIRONMENTAL PROTECTION

Do not contaminate water with the product or it's container. Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

### STORAGE AND DISPOSAL

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

DO NOT REUSE CONTAINER FOR ANY PURPOSE.

## IMPORTANT INFORMATION

FOR PROFESSIONAL USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose (litres of product/ha)	Maximum total dose (litres of product/ha/crop)	Latest time of application
Winter & spring wheat, winter & spring barley and Triticale	1,5	3	Before end of heading / inflorescence fully emerged (GS 59)

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

**Contents:  
10 litre**

170,00 mm

170,00 mm

## SAFETY PRECAUTIONS

### OPERATOR PROTECTION

**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES** when handling the concentrate or handling contaminated surfaces.

**WASH CONCENTRATE** from skin or eyes immediately.

**WASH HANDS AND EXPOSED SKIN** before eating or drinking and after work. IF YOU FEEL UNWELL, seek medical advice immediately (show label where possible).

### ENVIRONMENTAL PROTECTION

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### STORAGE AND DISPOSAL

**KEEP IN ORIGINAL CONTAINER**, tightly closed, in a safe place.

**DO NOT REUSE CONTAINER FOR ANY PURPOSE**

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

### RESISTANCE

KINGMAN contains folpet, a protectant fungicide from the phthalimide chemical group with multi-site contact activity.

### DISEASE CONTROL IN WHEAT, BARLEY AND TRITICALE

For the reduction of leaf spot (*Zymoseptoria tritici*) in winter and spring wheat and triticale and leaf blotch (*Rhynchosporium secalis*) in winter and spring barley and triticale. KINGMAN may be applied in a programme of 2 suitably timed sprays.

### DOSE RATE

1.5 L/ha per application

### SEPTORIA LEAF SPOT (*ZYMOSEPTORIA TRITICI*) IN WINTER AND SPRING WHEAT AND TRITICALE

As KINGMAN is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new growth from disease and therefore needs to be applied prior to further inspection. This is normally 3 weeks after the first application, but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or reinfection has occurred prior to application, KINGMAN should be mixed with alternative chemistry, e.g. prothioconazole, to provide curative activity (see Compatibility section).

### LEAF BLOTCH (*RHYNCHOSPORIUM SECALIS*) IN WINTER AND SPRING BARLEY AND TRITICALE

As KINGMAN is a protectant fungicide with contact activity, it is important to apply the first spray before the disease becomes established in the crop (normally GS 30-33). A second spray should be timed to protect new

growth from disease infection. This is normally 3 weeks after the first application, but may be sooner under conditions of rapid growth and high disease pressure. Where disease infection or reinfection has occurred prior to application, KINGMAN should be mixed with alternative chemistry, e.g. prothioconazole, to provide curative activity (see Compatibility section).

### WATER VOLUME

Apply KINGMAN in a water volume of 200-400 L/ha. AS KINGMAN has contact activity, it is important to achieve a good coverage of the crop when spraying.

### MIXING AND SPRAYING

For use by tractor mounted/trailer sprayer only. Before spraying it is important to check all hoses, filters and nozzles, and to ensure that the sprayer is clean and correctly calibrated to give an accurate application at the correct volume. Half-fill sprayer tank with clean water. Add the required amount of KINGMAN to the water and commence agitation. On emptying the container, RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely. Add the remainder of the water whilst continuing to agitate the mixture until spraying is complete.

Apply KINGMAN as a medium spray (as defined by the BCPC) in cereals. Thoroughly clean the sprayer and measuring equipment after using KINGMAN.

Crops should not be re-entered until spray residues are dry.

### COMPATIBILITY

Please consult DHM Agrochemicals Ltd. or your distributor for details on these products which may be mixed with KINGMAN in Ireland.

Maintain agitation during the mixing process and continue until the spraying is complete. Do not leave in spray tank for long period without using.

### Caution

After spraying a herbicide, sprayer tanks and equipment should be thoroughly cleaned before using KINGMAN.

### DISCLAIMER/CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

170,00 mm